

RESEARCH ARTICLE

CALIBRATION CURVE OF CEFADROXIL IN DIFFERENT SOLVENTS

Choundikar M. Y.*, Nitave S.A., Katageri A. R., Magdum S. V.

Dr. J. J. Magdum Trust's, Anil Alias Pintu Magdum Memorial Pharmacy College Dharnagutti.

Corresponding author: Choundikar M. Y


ABSTRACT:

In analytical chemistry, a calibration curve is a general method for determining the concentration of a substance in an unknown sample by comparing the unknown to a set of standard samples of known concentration. Chemical methods are developed to fit particular business, regulatory and research purposes. The objective of this research paper is to set out principles which should be followed when documenting methods of chemical testing. The calibration curve of Cefadroxil was plotted by using different solvents like distilled water, methanol, ethanol and dilute sodium hydroxide. Coefficient correlation (R^2) of Cefadroxil was found to be 0.999 in dil. sodium hydroxide.

Key words: Calibration curve, Cefadroxil, Coefficient correlation.

INTRODUCTION

The calibration curve is a plot of how the instrumental response, the so-called analytical signal, changes with the concentration of the analyte (the substance to be measured). The operator prepares a series of standards across a range of concentrations near the expected concentration of analyte in the unknown. [1] The concentrations of the standards must lie within the working range of the technique (instrumentation) they are using (figure01). Analyzing each of these standards using the chosen technique will produce a series of measurements. For most analyses a plot of instrument response vs. analyte concentration will show a linear relationship. The operator can measure the response of the unknown and, using the calibration curve, can interpolate to find the concentration of analyte. [2, 3, 4]

Figure1: Calibration curve

A general method for analysis of concentration involves the creation of a calibration curve. This allows for determination of the amount of a chemical in a material by comparing the results of unknown sample to those of a series known standards. If the concentration of element or compound

in a sample is too high for the detection range of the technique, it can simply be diluted in a pure solvent. If the amount in the sample is below an instrument's range of measurement, the method of addition can be used. In this method a known quantity of the element or compound under study is added, and the difference between the concentration added, and the concentration observed is the amount actually in the sample. [5]

Cefadroxil is a broad-spectrum antibiotic of the cephalosporin type, effective in Gram-positive and Gram-negative bacterial infections. It is a bactericidal antibiotic. Cefadroxil, (6*R*,7*R*)-7-[[*(2R)*-2-amino-2-(4-hydroxyphenyl)acetyl]amino]-3-methyl-8-oxo-5-thia-1-azabicyclo[2.2.1]heptane-2-carboxylic acid is a first-generation cephalosporin antibacterial drug that is the para-hydroxy derivative of cefalexin, and is used similarly in the treatment of mild to moderate susceptible infections such as the bacterium *Streptococcus pyogenes*, causing the disease popularly called strep throat or streptococcal tonsillitis, urinary tract infection, reproductive tract infection, and skin infections.




Figure 2: Structure of Cefadroxil

Molecular Formula: $C_{16}H_{17}N_3O_5S, H_2O$

Molecular Weight: 381.40

Category: Antibacterial.

Storage: Store in tightly-closed containers in a cool place.

MATERIALS AND METHODS:

Materials:

Cefadroxil was procured as a gift sample from Sunrise pharmaceutical Ltd Mahisal, Maharashtra and all other chemicals of AR grade were obtained from Research lab fine Mumbai Maharashtra, India. The instrument used were JASCO- V- 550, UV/VIS double beam spectrophotometer, Digital electronic balance (Shimadzu-Japan).

Solubility and solvent effect:

A most suitable solvent is one that does not itself absorb in the region under investigation. Most commonly used solvent is 95% ethanol. Ethanol is a best solvent as it is cheap and is transparent down to 210 m μ . hexane and other hydrocarbons can be used because these are less polar and have least interaction with the molecule under investigation.

Table No. 01 Solubility of Cefadroxil in different solvents

Solvent	Solubility
Distilled Water	soluble
Dil. NaOH	Slightly soluble
Ethanol	Slightly soluble
Methanol	Slightly soluble

Choice of solvent:

1. It should not itself absorb radiations in the region under investigation.
2. It should be less polar so that it has minimum interaction with the solute molecules.

Calibration curve of Cefadroxil in different solvents: [8]**1. Cefadroxil in Distilled Water:**

Drug : Cefadroxil
Solvent: Distilled Water

Procedure:

100 mg Cefadroxil was weighed accurately and dissolved in 100 ml of distilled water in volumetric flask. Flask was shaken for 5 minutes to dissolve drug properly. Flask was labeled as Stock Solution. 1 ml of stock solution was further diluted into 100 ml of distilled water. max was determined by scanning on UV – Visible spectrophotometer. Further dilutions were prepared by diluting 1 ml stock solution in 100 ml, 2ml stock solution in 100 ml, and so on.

2. Cefadroxil in dil. NaOH:

Drug : Cefadroxil
Solvent: dil. NaOH

Procedure:

100 mg Cefadroxil was weighed accurately and dissolved in 100 ml of dil. NaOH in volumetric flask. Flask was shaken for 5 minutes to dissolve drug properly. Flask was labeled as Stock Solution. 1 ml of stock solution was further diluted into 100 ml of dil. NaOH. max was determined by scanning on UV – Visible spectrophotometer. Further dilutions were prepared by diluting 1 ml stock solution in 100 ml, 2ml stock solution in 100 ml, and so on.

Cefadroxil in Ethanol:

Drug : Cefadroxil
Solvent: Ethanol

PROCEDURE:

100 mg Cefadroxil was weighed accurately and dissolved in 100 ml of ethanol in volumetric flask. Flask was shaken for 5 minutes to dissolve drug properly. Flask was labeled as Stock Solution. 1 ml of stock solution was further diluted into 100 ml of ethanol. max was determined by scanning on UV – Visible spectrophotometer. Further dilutions were prepared by diluting 1 ml stock solution in 100 ml, 2ml stock solution in 100 ml, and so on.

4 Cefadroxil in Methanol:

Drug : Cefadroxil
Solvent: Methanol

Procedure:

100 mg Cefadroxil was weighed accurately and dissolved in 100 ml of methanol in volumetric flask. Flask was shaken for 5 minutes to dissolve drug properly. Flask was labeled as

Stock Solution. 1 ml of stock solution was further diluted into 100 ml of methanol. max was determined by scanning on UV – Visible spectrophotometer. Further dilutions were prepared by diluting 1 ml stock solution in 100 ml, 2ml stock solution in 100 ml, and so on

RESULT AND DISCUSSION:

Cefadroxil in distilled water:

Figure 3: Scanned max value 239





Figure 4: Calibration curve of Cefadroxil in Distilled water.



Cefadroxil in dil. NaOH:

Figure 5: Scanned max value 238







Figure: 6 Calibration curve of Cefadroxil in dil.NaOH**Cefadroxil in Ethanol**

Scanned max value : 234.2 nm.

Figure: 7 Calibration curve of Cefadroxil in ethanol**Cefadroxil in Methanol:**

Scanned max value : 230 nm.

Figure: 8 Calibration curve of Cefadroxil in Methanol

CONCLUSION:

The calibration curve provides a reliable way to calculate the uncertainty of the concentration calculated from the calibration curve (using the statistics of the least squares line fit to the data). [5] The calibration curve provides data on an empirical relationship. The mechanism for the instrument's response to the analyte may be predicted or understood according to some theoretical model, but most such models have limited value for real samples. (Instrumental response is usually highly dependent on the condition of the analyte, solvents used and impurities it may contain; it could also be affected by external factors such as pressure and temperature). [6]

Calibration curve of Cefadroxil was performed in different solvents viz. distilled water, dil. Sodium hydroxide, ethanol and methanol. Coefficient correlation (R^2) of Cefadroxil was found to be 0.999 in dilute sodium hydroxide and less than 0.999 in methanol, distilled water and ethanol. Hence this result can further be used in various research related to Cefadroxil.

REFERENCES:

1. Sharma BK. In; Instrumental Methods of Chemical Analysis, 20th Edn., Krishna Prakashan Media, Ltd., Meerut, 2001:4-6.
2. Hollas JM. In; Modern Spectroscopy, 4th Edn., John Wiley and Sons Ltd., Chichester, 2004: XIX.
3. Conners KA. A Textbook of Pharmaceutical Analysis, 3rd Edn., Wiley-Interscience Publication, John Wiley & Sons, 1982:173-187.
4. R.M. Silvesterin and G.C. Bassler. Spectrometric Identification of Organic Compounds.
5. Meiling Qi, Peng Wang, YujingSun, Jun Wang. J Liq Chrom Rel Technol 2003; 26:1927-1936.
6. Thorburn Burns D, O'Callaghan M, Franklin Smyth W, Ayling C. J. Fresenius. J Anal Chem 1991; 340:53-56.
7. Hsu Mei-Chich, Cheng Mei-Chai. J Chrom A 1991; 549:410-415.
8. International Conference on Harmonization, Draft Guideline on Validation Procedure, Definition and Terminology Federal Register, 1995, 60, 11260