

Original Research Article

DIFFERENT FRACTIONS OF ANTIOXIDANT ACTIVITY OF SOME MEDICINAL PLANT

Baburao Bhukya*

Nethaji Institute of Pharmaceutical Sciences, Somidi, Kazipet, Warnagal, Telangana State-506003.

Corresponding Author: Dr. B BABU RAO, M.Pharm., Ph.D. Professor and Principal, Nethaji Institute of Pharmaceutical Sciences, Somidi, Kazipet, Warnagal, Telangana State-506003.

ABSTRACT:

The present study was carried out to evaluate the free radical scavenging activity of various fractions of *Hiptage benghalensis* (Linn) Kurz. DPPH free radical and ABTS free radical scavenging assays were carried out to evaluate the antioxidant potential of the extract. The antioxidant activity of toluene, ethyl acetate, butan-2-one and n-butyl alcoholic extract increased in dose depending manner. In DPPH radical scavenging assay the IC₅₀ value for DPPH scavenging by the toluene, ethyl acetate, butan-2-one and n-butyl alcoholic extracts of *Hiptage benghalensis* were 31.3, 15.4, 29.4 and 33.2 respectively. The IC₅₀ value for ABTS free radical scavenging ability of toluene, ethyl acetate, butan-2-one and n-butyl alcoholic extracts of *Hiptage benghalensis* were 6.25, 1.27, 9.71 and 7.83 respectively. The results obtained in this study indicate that the ethyl acetate extract can be potential source of natural antioxidant.

Key Words: *Hiptage benghalensis*, antioxidant, DPPH radical, ABTS free radical.

INTRODUCTION

Natural products remain a prolific source for the discovery of new drugs and drugs leads even from Vedic period. Recent data suggests that 80% of drug molecules are natural products or natural compound inspired. Studies on sources of new drugs 1981 to 2007 reveal that almost half of the drugs approved since 1994 are based on natural products (Butler, 2008). Indian natural products, particularly those from traditional medicinal plants which are reported in the classic texts like Ayurveda and Charak Samhita, have contributed towards drug discovery.

In India the plant is widely distributed in forest at an altitude of 1800 m, except in dry regions. And also found in moist habitats such as along riverbanks and in ravines. The plant occurs in Sri Lanka, Andaman Islands, Bangladesh, and Myanmar (Burma) to Southern China. The bark, leaves and flowers are aromatic in nature and are commonly used as refrigerant, expectorant, cardiogenic, Anti-inflammatory and insecticidal. They are used in burning sensation, wounds, ulcers, leprosy, cardiac debility, rheumatism and hyperdipsia (Chatterjee and Chandrapakrashi, 1994; Vaidyanam, 1995). In Ayurveda the leaves and barks are considered vulnerary, the leaves are highly regarded for treating

skin diseases. The leaf juice possesses insecticidal properties and is used an external application for scabies. The plant is also used in the treatment of chronic rheumatism and asthma (Parrotta, 2001).

MATERIALS AND METHODS

Plant material:

The leaves of *Hiptage benghalensis* (Linn) Kurz were collected in the month of June from thirumal hills, Andhra Pradesh, India. The selected plant was authenticated by Prof. Raju S. Vastavaya, Department of Botany, Kakatiya University, Warangal and voucher specimens were being maintained in the herbarium of University College of Pharmaceutical Sciences, Kakatiya University, Warangal.

Preparation of Extracts:

The leaves of *Hiptage benghalensis* (3kg) were made free from the adherent foreign material and air-dried. Then they were coarsely powdered and 2kg of each was macerated with methanol in a round bottom flask for 7 days separately. The content of the flask were stirred intermittently to ensure the efficiency of the extraction. After a week, they were filtered and concentrated under reduced pressure to yield corresponding extracts, and the extracts were kept in a desiccator to remove moisture and stored properly until reuse.

The methanolic extracts of *Hiptage benghalensis* were dispersed in sufficient amount of distilled water separately and fractionated with toluene, ethyl acetate, butan-2-one and n-butyl alcohol in succession. The obtained fractions and the aqueous residues were concentrated under reduced pressure to yield corresponding extracts.

Antioxidant Activity

DPPH free radical scavenging effect (Blois, 1958)

The free radical scavenging activity of MHB fractions were measured by DPPH using the method of Blois. Ascorbic acid was used as a reference standard. The methanolic solution of DPPH (0.2mM) was added to different concentrations (100, 200, 400, 600, 800, 1000 µg/ml in methanol) of fractions. After 30 min, absorbance was measured at 517 nm. The degree of discoloration indicates the scavenging potential of the extract. Lower absorbance of the reaction mixture indicates higher free radical scavenging activity. The IC₅₀ (Inhibitory Concentration) is the concentration of sample required to scavenge 50% of DPPH free radicals.

ABTS free radical scavenging effect (Re et al., 1999)

ABTS was dissolved in water to a 7mM concentration. ABTS radical cation (ABTS•1) was produced by reacting ABTS stock solution with 2.45 mM potassium persulfate (final concentration) and allowing the mixture to stand in the dark at room temperature for 12–

16 h before use . Because ABTS and potassium persulfate react stoichiometrically at a ratio of 1:0.5, this will result in incomplete oxidation of the ABTS. Oxidation of the ABTS commenced immediately, but the absorbance was not maximal and stable until more than 6 h had elapsed. The radical was stable in this form for more than two days when stored in the dark at room temperature. For the study of antioxidant activity, various concentrations (100, 200, 300, 400, 500µg/ml) of extracts were added to the ABTS•1 solution. Then the mixture was diluted with ethanol to get an absorbance of 0.70 to 0.80 at 734 nm and equilibrated at 30°C. Stock solutions. After addition of 1.0ml of diluted ABTS•1 solution appropriate solvent blanks were run in each assay.

Calculation :


$$\text{Percentage O.D.} = \frac{\text{Control O.D.} - \text{Test O.D.}}{\text{Control O.D.}} \times 100$$

RESULTS AND DISCUSSION:

In Vitro DPPH free radical scavenging activity of Selected Fractions MHB Table 1.

Compound	IC50 (µg/ml)
STD	4.3
T-MHB	31.3
EA-MHB	15.4
BN-MHB	29.4
BL-MHB	33.2


Fig: 1 *In Vitro* DPPH free radical scavenging activity of Selected Fractions MHB



In Vitro ABTS free radical scavenging activity of Selected Fractions MHB Table 2

Compound	IC ₅₀ (µg/ml)
Ascorbic acid	2.42
T-MHB	6.25
EA-MHB	1.27
BN-MHB	9.7
BL-MHB	7.83

Fig: 2 *In Vitro* ABTS free radical scavenging activity of Selected Fractions MHB



The reduction capability of the DPPH radical induced by antioxidants is determined by the decrease in the absorbance at 517 nm. The decrease in the absorbance of DPPH radical caused by antioxidants, due to the reaction between antioxidant molecules and radical, progresses which results in the scavenging of the radical by hydrogen donation. It is visually noticeable as a change in color from purple to yellow. All the fractions of MHB were evaluated for their free radical activity against DPPH radicals and the results were expressed as IC₅₀ values and were shown in Table 1 & Fig 1. This indicated the antioxidant activity of all fractions against DPPH free radical. Among all the fractions showed a maximum scavenging potential with IC₅₀ value of 31.3, 15.4, 29.4 and 33.2 µg/ml. Among all fractions of MHB, the antioxidant potentials was found as EA-MHB > T-MHB > BN-MHB > BL-MHB.

It is a method for the screening of antioxidant activity. It reported as a decolorization assay applicable to both lipophilic and hydrophilic antioxidants, including flavonoids, hydroxycinnamates, carotenoids, and plasma antioxidants. The pre-formed radical monocation of 2, 2'-azinobis-(3-ethylbenzothiazoline-6-sulfonic acid) (ABTS•1) is generated by oxidation of ABTS with potassium persulfate and is reduced in the presence of such hydrogen-donating antioxidants. The influences of both the concentration of antioxidant and duration of reaction on the inhibition of the radical cation absorption are taken into account when determining the antioxidant activity. All

the fractions of MHB were evaluated for their free radical activity against ABTS radicals and the results were expressed as IC₅₀ values and were shown in Table 2 & Fig 2. This indicated the antioxidant activity of all fractions against ABTS free radical. Among all the fractions showed a maximum scavenging potential with IC₅₀ value of 6.25, 1.25, 9.5 and 7.83µg/ml. The radical scavenging activity of MHB fractions decreased in the following order: EA-MHB> T-MHB> BL-MHB> BN-MHB.

This information encouraged us to evaluate the free radical scavenging effect of MHB fractions by employing widely used methods i.e. DPPH free radical scavenging activity, ABTS assay, radical scavenging activity. MHB fractions showed moderate to strong free radical scavenging activities. These results confirmed that the fractions of the selected plant exhibited a significant antioxidant activity.

REFERENCES

1. Butler M S, Natural Products to drug; natural product-derived compounds in clinical trials, Nat Prod Rep 2008.25: 475.
2. Chatterjee A, S. Chandra Pakrashi, The treatise on Indian medicinal plants, New Delhi, 1994. Vol.3: 130-132.
3. Kirtikar K R, Basu B D, Indian Medicinal Plants, second edition, International Book Distributors, Deharadun, 1988. 3: 2017-18.
4. John a Parrott, Healing medicinal plants of peninsular India. CAB International publishing New York, 2001. pp 614-615.
5. Blois M S, Antioxidant determinations by the use of a stable free radical. Nature 1958. 29: 1199-2000.
6. Re R, Pellegrini N, Proteggente A, Pananath A, Yang M, Rice-Evans C, Free Radical Biology and Medicine, 1999. 26: 1231-1237.
7. Vaidyanam P S, Indian Medicinal Plants a compendium of 500 Species, Orient Longman limited. 1995. Vol-3:153.